

A&A manuscript no.
(will be inserted by hand later)

Your thesaurus codes are:
04 (11.09.1 NGC 4418; 11.09.4; 11.19.1; 13.09.4)

ASTRONOMY
AND
ASTROPHYSICS

The obscured mid-infrared continuum of NGC 4418: a dust- and ice-enshrouded AGN ^{*}

H.W.W. Spoon¹, J.V. Keane², A.G.G.M. Tielens², D. Lutz³, and A.F.M. Moorwood¹

¹ European Southern Observatory, Karl-Schwarzschild Strasse 2, D-85748 Garching, Germany
email: hspoon@eso.org, amoor@eso.org

² Kapteyn Astronomical Institute, Postbus 800, NL-9700 AV Groningen, The Netherlands
email: j.keane@astro.rug.nl, tielens@astro.rug.nl

³ Max-Planck-Institut für Extraterrestrische Physik (MPE), Postfach 1312, D-85741 Garching, Germany
email: lutz@mpe.mpg.de

Abstract. We report the detection of absorption features in the 6–8 μm region superimposed on a featureless mid-infrared continuum in NGC 4418. For several of these features this is the first detection in an external galaxy. We compare the absorption spectrum of NGC 4418 to that of embedded massive protostars and the Galactic centre, and attribute the absorption features to ice grains and to hydrogenated amorphous carbon grains. From the depth of the ice features, the powerful central source responsible for the mid-infrared emission must be deeply enshrouded. Since this emission is warm and originates in a compact region, an AGN must be hiding in the nucleus of NGC 4418.

Key words: Galaxies: individual: NGC4418 — Galaxies: ISM — Galaxies: Seyfert — Infrared: ISM: lines and bands

1. Introduction

The ISO mission has considerably enhanced our knowledge of the mid-IR properties of normal, starburst, Seyfert and Ultra-luminous Infrared Galaxies (ULIRGs). The spectra of most sources are dominated by ISM emission features, the most prominent of which are the well-known PAH emission bands at 6.2, 7.7, 8.6, 11.3 and 12.7 μm . The PAH features and the emission lines have been used qualitatively and quantitatively as diagnostics for the ultimate physical processes powering the galactic nuclei (Genzel et al. 1998; Lutz et al. 1998; Rigopoulou et al. 1999; Tran et al. 2000). A broad absorption band due to the Si-O stretching mode in amorphous silicates, centered at 9.7 μm , is also commonly detected in galaxies. Since the center of the silicate absorption coincides with a gap between the 6.2–8.6 μm and 11.3–12.8 μm PAH complexes,

it is not readily apparent whether a 9.7 μm flux minimum should be interpreted as the “trough” between PAH emission features or as strong silicate absorption, or as a combination of the two. In spectra observed towards heavily extincted Galactic lines of sight, a strong silicate feature is often accompanied by ice absorption features in the 6–8 μm region (e.g. Whittet et al. 1996). Until recently this combination had not been reported in equally extincted extragalactic sources, despite detections of ice features at shorter wavelengths (Spoon et al. 2000; Sturm et al. 2000). In this *Letter* we report on the detection of ices in the strongly absorbed, ISO-PHT-S spectrum of NGC 4418, a nearby ($D=29$ Mpc; $1''=140\text{pc}$) luminous ($L_{\text{IR}}=10^{11} L_{\odot}$) bright IRAS galaxy. NGC 4418 and the distant ULIRG IRAS 00183-7111 (Tran et al. 2000) are the first detections of these ice features in external galaxies.

NGC 4418 is well-known for its deep 9.7 μm silicate feature (Roche et al. 1986). Additional evidence for strong extinction is the weakness ($H\alpha$) and absence ($H\beta$, $\text{Br}\alpha$, $\text{Br}\gamma$) of hydrogen recombination line emission (Kawara et al. 1989; Ridgway et al. 1994; Lehnert & Heckman 1995; L.Kewley, priv.comm.), commonly detected in galaxies. HST-NICMOS images (Scoville et al. 2000) show hardly any structure in the inner $400\text{pc}\times 400\text{pc}$, except for large scale extinction ($\Delta A_V \sim 2$). The IRAS colors of NGC 4418 indicate that — unlike most other galaxies — the 12–100 μm emission of NGC 4418 is dominated by a warm dust component, peaking shortward of 60 μm (see Fig. 1). VLA radio maps at 6 and 20 cm (Condon et al. 1990; Eales et al. 1990) show NGC 4418 to be compact ($70\text{pc}\times 50\text{pc}$ at most). This, as well as the presence of large quantities of warm dust, has been taken as evidence for the presence of an otherwise hidden AGN in NGC 4418. In this *Letter* we present mid-IR spectral evidence lending further support for the presence of an AGN in NGC 4418.

2. Observations

A low resolution ($\lambda/\Delta\lambda \sim 90$) ISO-PHT-S spectrum of the central $24''\times 24''$ of NGC 4418 was obtained on 1996

Send offprint requests to: H.W.W. Spoon

^{*} Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) and with the participation of ISAS and NASA




Fig. 1. The 1-3000 μm spectral energy distribution of NGC 4418, including the new ISO-PHT-S, ISO-CAM and smoothed ISO-LWS data. The data shown in grey is the spectrophotometry of Roche et al. (1986). Other data shown has been taken from Dudley (1997), Lisenfeld et al. (2000), Roche & Chandler (1993), Soifer et al. (1989) and Wynn-Williams & Becklin (1993). The dotted and dashed curves show two choices for the local mid-IR continuum. The dotted line interpolates the peaks of the observed data, whereas the dashed line assumes a stronger 18 μm silicate absorption.

July 14 as part of a project on the interstellar medium of normal galaxies (Helou et al. 2000). The measurement was carried out in triangular chopped mode, using a chopper throw of 150". The resulting spectrum is thus free of contributions from zodiacal light. The ISO-PHT-S data were reduced using PIA 8.2. The absolute calibration is accurate to within 20%. The resulting spectrum is shown in Fig. 2.

In Fig. 1 we compile the 1-3000 μm (spectro)photometric observations of NGC 4418, including the new, standard reduced, ISO-PHT-S, ISO-CAM and ISO-LWS observations. Stellar emission probably dominates the near-IR regime up to 4 μm , beyond which a strong mid-infrared continuum sets in. The SED reveals that a substantial part of the emission emanates from optically thick warm dust (Roche & Chandler 1993; Lisenfeld et al. 2000), peaking at 40-60 μm . Also shown in Fig. 1 are two equally possible choices of local continuum which we have adopted to analyse the ice, hydrogenated amorphous carbon (HAC) and silicate absorption. This will be discussed in detail in Section 3.2.

3. The mid-IR spectrum of NGC 4418

3.1. Spectral features

The mid-IR spectrum of NGC 4418 (Fig. 2) bears little resemblance to the spectrum of (almost) any other




Fig. 2. A comparison of the ISO spectra of NGC 4418, M 82, the embedded massive protostar W 33A and the Galactic centre (Sgr A*). Except for NGC 4418, the spectra have been scaled and offset. We removed the strong 7 μm [Ar II] line from the Sgr A* spectrum. The vertical dashed lines facilitate comparison between the four spectra with well known Galactic absorption features. The arrows mark the rest wavelengths of the 6.2, 7.7, 8.6 and 11.3 μm PAH features. The zero flux levels for M 82, NGC 4418 and W 33A are indicated with horizontal dotted lines.

galaxy obtained by ISO. The spectra of normal and starburst galaxies (Rigopoulou et al. 1999; Helou et al. 2000) are dominated by strong PAH emission features. Seyfert galaxies with a clear line of sight to AGN-heated hot dust, on the other hand, are dominated by a strong mid-IR continuum with PAHs barely recognizable (Clavel et al. 2000). The mid-IR spectrum of NGC 4418 is rich in features but does not resemble PAH spectra. Rather it is similar to spectra observed towards heavily extinguished Galactic lines of sight, such as deeply embedded massive protostars or the Galactic centre (Fig. 2). These show no evidence for PAH emission features but do show strong absorption features. A simple criterion for the role of PAH emission and absorption features is based on the location of maxima in the 6-7 μm region: PAH spectra show the 6.2 μm emission feature, whereas absorption spectra show a peak at 6.5-6.7 μm , which is not an emission feature but a window of reduced absorption between two features.

A detailed comparison of the absorption features with Galactic templates may be able to shed further light on the origin of the extinction in NGC 4418. The absorption features were determined by fitting a local polynomial to the peaks of the 5.8-8 μm spectrum of NGC 4418. The absence of the 6.2 μm PAH feature indicates that emission features are very weak in the spectrum, and justifies this simple procedure to derive the shape and depth of the absorption features. Fig. 3 shows the resulting NGC 4418

Table 1. Observed parameters for the NGC 4418 features.

λ [μm]	τ	τ_{int} [cm^{-1}]	carrier	N [10^{17} cm^{-2}]
6.0	0.8	87	H ₂ O	73
6.85	1.1	27	HAC?	~ 200
7.3	0.6	12	HAC?	~ 200
7.67	0.12	1.2	CH ₄	1.6
9.7	7	—	silicates	—
18	~ 1.5	—	silicates	—
			atomic H	$\sim 2 \times 10^6$ *

* Column to the mid-IR dust source calculated assuming Galactic gas to dust ratios. Note that the X-ray absorbing column to the central engine may be even higher.

optical depth spectrum along with the optical depth spectrum of ices towards the massive protostar W 33A (Gibb et al. 2000) and the spectrum of the Galactic centre (Sgr A*), which displays absorptions due to ices as well as features due to dust in the diffuse ISM (Chiar et al. 2000). The spectrum of NGC 4418 shows absorption features at 6.0, 6.8, 7.3, 7.6, 10 and 18 μm (Table 1). The 6.0 μm feature in NGC 4418 is similar to that in the Galactic sources but with a perhaps more pronounced long wavelength wing. The 6.85 μm feature is considerably narrower than the molecular cloud feature but is similar in width to the diffuse ISM feature. The band at 7.3 μm is substantially broader than the molecular cloud and diffuse ISM features. The absorption band near 7.6 μm is similar to that observed locally.

The presence of ice along the line of sight toward NGC 4418 is suggested by the identification of the 7.6 μm band with CH₄ (Boogert et al. 1996, 1997) and by the presence of the 6.0 μm band due to H₂O ice (Keane et al. 2000; Chiar et al. 2000). The origins of the 6.85 μm and 7.3 μm bands in NGC 4418 are unclear. Interstellar ices also show features at these wavelengths, however, their relative strengths as well as widths are markedly different in NGC 4418. The 6.85 μm and 7.3 μm band ratios are consistent with the features observed towards Sgr A*, which have been attributed to CH deformation modes in HAC-like dust grains (Chiar et al. 2000). Thus, as for the Galactic centre, both ice characteristics for shielded dense molecular cloud environments and HAC-like grain characteristics for diffuse ISM dust seem to be present along the line of sight. This conclusion could be tested through observations in the 3 μm window, which contains the strong 3 μm H₂O ice band and the 3.4 μm CH stretching modes of HAC materials. Guided by variations seen for the Galactic centre region (Chiar et al. 2000), and between M 82 and NGC 1068 (Sturm et al. 2000), we speculate that the relative weight of ice and HAC components may vary considerably among galaxies.




Fig. 3. The optical depth spectra for W 33A, NGC 4418, and Sgr A*. The vertical lines indicate the positions of the 6.0, 6.85, 7.3, and 7.6 μm ice and HAC absorption bands seen toward Galactic lines of sight.

3.2. Dust and ice column densities

The column densities of the ice species can be derived by dividing the integrated optical depth (τ_{int}) by the molecular band strength (Gerakines et al. 1995; Boogert et al. 1997). Table 1 summarizes the computed column densities of CH₄ and H₂O ice. The column densities are consistent with those derived for embedded massive protostellar objects in molecular clouds (Boogert et al. 1996; Keane et al. 2000). Also shown in Table 1 are the calculated HAC column densities for the 6.85 and 7.3 μm bands, assuming the intrinsic integrated band intensities for saturated aliphatic hydrocarbons from values by Wexler (1967). These integrated intensities are stronger than other current literature values (Sandford et al. 1991; Furton et al. 1999). For NGC 4418 a substantial fraction of the carbon is locked up in HAC ($\sim 20\%$) as compared to the Galactic centre (a few % in Sgr A*; Pendleton et al. 1994).

As mentioned previously (Sect. 2), two continua, differing only longward of 8 μm were considered for NGC 4418 (Fig. 1). The effect of the different continua on the silicate feature is noticeable when fitting the silicate profile by the Galactic Centre Source GCS 3, a pure absorption feature, i.e. no intrinsic emission (Figer et al. 1999). However, the exact optical depth is still difficult to determine due to saturation of the silicate feature in NGC 4418. Adopting the dotted line in Figure 1 results in a NGC 4418 silicate profile in which the blue wing is well matched but the red wavelength wing is poorly fitted by the GCS 3 profile. If however the dashed line is adopted, then the fit to the red wing improves while still maintaining a reasonable match to the blue wing of the 10 μm silicate band. The 9.7 μm and 18 μm optical depths for the second case are $\tau_{9.7} \sim 7$ (corresponding to $A_V \sim 130$) and $\tau_{18} \sim 1.5$. These numbers, along with the derived hydrogen column density,

are shown in Table 1. The apparent optical depth ratio of the $9.7/18\mu\text{m}$ silicate bands, regardless of adopted continuum, is significantly larger than the ratio determined by Demyk et al. (1999) for two Galactic protostars. This might suggest that complex radiative transfer effects are important, which are however beyond the scope of this Letter.

4. Discussion and conclusions

We have compared the ISO-PHT-S spectrum of NGC 4418 with spectra of our template sources and found no sign of PAH emission, neither from the nucleus, nor from that part of the disk contained within the $24'' \times 24''$ ISO-PHT-S aperture. Instead we found deep absorption features imposed upon a featureless mid-IR continuum. We identify the $6\text{--}8\mu\text{m}$ absorption features with foreground ices and HAC-like grains.

The nature of the central source in NGC 4418 cannot be inferred from the observed mid- to far-IR spectrum alone, given the absence of any “signposts”, like the $6.2\mu\text{m}$ PAH emission feature ($F_{\text{PAH}} < 6 \times 10^{-20} \text{ W/cm}^2$), and of fine structure lines ([O I], [C II] and [O III], Malhotra et al. (1999); $F_{[\text{Ne II}]} < 2.3 \times 10^{-20} \text{ W/cm}^2$, from archival ISO-SWS data). Both a heavily enshrouded AGN or a similarly extinguished nuclear starburst could be responsible for the observed continuum. Even if a starburst could be accommodated within the compact central source ($< 70 \text{ pc}$ in the mid-IR (Scoville et al. 2000) and $25\text{--}70 \text{ pc}$ at 6 and 20 cm (Eales et al. 1990; Kawara et al. 1990)), it would be highly unlikely to block the escape of any mid-IR starburst indicator from a region of that size. The most likely origin is therefore a heavily enshrouded AGN as suggested previously by Roche et al. (1986), Kawara et al. (1990) and Dudley & Wynn-Williams (1997). The $0.1''$ point source (5 mJy) detected with the Parkes Tidbinbilla Interferometer (PTI) at 13 cm (Kewley et al. 2000; L.Kewley, priv.comm.) may actually pinpoint the AGN itself. Far-IR to millimetre sizes are less well constrained but the warm IRAS colours suggest that the emission in this range also arises in the nuclear region.

Our finding of strong absorptions due to cold silicates and ices in NGC 4418 leads us to believe that the same absorptions may be present in the mid-IR spectra of other galaxies. Indeed, we have found similar absorptions in about a dozen of 225 galaxies observed spectroscopically by ISO. Spectral identifications have to be done with great care since simultaneous presence of PAH emission makes other spectra more complex than the one of NGC 4418.

Since the overall shape of the NGC 4418 $6\text{--}11\mu\text{m}$ spectrum with its maximum near $8\mu\text{m}$ mimicks at first glance a PAH spectrum, we point out the need for high S/N spectra to clearly identify the indicators for bona fide PAH spectra or absorption dominated spectra. The most obvious discriminator is the $6.2\mu\text{m}$ PAH peak to be contrasted with the $6.5\text{--}6.7\mu\text{m}$ pseudo-maximum in absorption spec-

tra, which is due to a window between two absorption features. In a forthcoming paper we will address this issue for our large ISO galaxy sample.

Acknowledgements. The authors wish to thank Olivier Laurent and Leticia Martín-Hernández for performing the ISO-CAM and ISO-LWS data reduction, respectively, as well as Ilse van Bemmelen, Peter Barthel, Matthew Lehnert, Dave Sanders, Nick Scoville, Ralf Siebenmorgen, Eckhard Sturm and Dan Tran for discussions.

References

- Boogert A.C.A., Schutte W.A., Tielens A.G.G.M., et al., 1996, A&A 315, L377
- Boogert A.C.A., Schutte W.A., Helmich F.P., et al., 1997, A&A 328, 649
- Chiar J.E., Tielens A.G.G.M., Whittet D.C.B., et al., 2000, ApJ 537, 749
- Clavel J., Schulz B., Altieri B., et al., 2000, A&A 357, 839
- Condon J.J., Helou G., Sanders D.B., Soifer B.T., 1990, ApJS 73, 359
- Demyk K., Jones A.P., Dartois E., et al., 1999, A&A 349, 267
- Dudley C.C., 1997, Ph.D. thesis, Univ. Hawaii
- Dudley C.C., Wynn-Williams C.G., 1997, ApJ 488, 720
- Eales S.A., Becklin E.E., Hodapp K.-W., et al., 1990, ApJ 365, 478
- Figer D.F., McLean I.S., Morris M., 1999, ApJ 514, 202
- Furton D.G., Laiho J.W., Witt A.N., 1999, ApJ 526, 752
- Genzel R., Lutz D., Sturm E., et al., 1998, ApJ 498, 579
- Gerakines P.A., Schutte W.A., Greenberg J.M., van Dishoeck E.F., 1995, A&A 296, 810
- Gibb E.L., Whittet D.C.B., Schutte W.A., et al., 2000, ApJ 536, 347
- Helou G., Lu N.Y., Werner M.W., Malhotra S., et al., 2000, ApJ 532, L21
- Kawara K., Nishida M., Phillips M.M., 1989, ApJ 337, 230
- Kawara K., Taniguchi Y., Nakai N., et al., 1990, ApJ 365, L1
- Keane J.V., Tielens A.G.G.M., Boogert A.C.A., et al., 2000, A&A, Submitted
- Kewley L.J., Heisler C.A., Dopita M.A., et al., 2000, ApJ 530, 704
- Lehnert M.D., Heckman T.M., 1995, ApJS 97, 89
- Lisenfeld U., Isaak K.G., Hills R., 2000, MNRAS 312, 433
- Lutz D., Spoon H.W.W., Rigopoulou D., et al., 1998, ApJ 505, L103
- Malhotra S., et al., 1999, In: The Universe as seen by ISO, P. Cox, M.F. Kessler (eds.), ESA-SP 427, 813
- Pendleton Y.J., Sandford S.A., Allamandola L.J., et al., 1994, ApJ 437, 683
- Ridgway S.E., Wynn-Williams C.G., 1994, 428, 609
- Rigopoulou D., Spoon H.W.W., Genzel R., et al., 1999, AJ 118, 2625
- Roche P.F., Aitken D.K., Smith C.H., James S.D., 1986, MNRAS 218, 19P
- Roche P.F., Chandler C.J., 1993, MNRAS 265, 486
- Sandford S.A., Allamandola L.J., Tielens A.G.G.M., et al., 1991, ApJ 371, 607
- Scoville N.Z., Evans A.S., Thompson R., et al., 2000, AJ 119, 991
- Soifer T., Böhrer L., Neugebauer G., et al., 1989, AJ 98, 766

- Spoon H.W.W., Koornneef J., Moorwood A.F.M., et al., 2000, A&A 357, 898
- Sturm E., Lutz D., Tran D., et al., 2000, A&A 358, 481
- Tran Q.D., Lutz D., Genzel R., et al., ApJ, Submitted
- Wexler A.S., 1967, Applied Spec. Rev. 1, 29
- Whittet D.C.B., Schutte W.A., Tielens A.G.G.M., et al., 1996, A&A 315, L357
- Wynn-Williams C.G., Becklin E.E., 1993, ApJ 412, 535